





MAY 5 1964

LIBRARY  
OF THE  
UNIVERSITY  
OF ILLINOIS

590.5

FI

v. 34

BIOLOGY.

Return this book on or before the  
**Latest Date** stamped below.

Theft, mutilation, and underlining of books  
are reasons for disciplinary action and may  
result in dismissal from the University.

University of Illinois Library

JAN 12 1966

DEC 4 1967

SEP 15 1971

~~NOV 14 1988~~

~~DECLERED~~

DEC 14 1990

DEC 07 1992







34<sup>18</sup>

# FIELDIANA • ZOOLOGY

Published by  
CHICAGO NATURAL HISTORY MUSEUM

Volume 34

JULY 24, 1953

No. 18

## A COLOMBIAN RACE OF *TINAMUS OSGOODI*

EMMET R. BLAKE

ASSOCIATE CURATOR, DIVISION OF BIRDS

1953

A small collection of game birds made in Colombia by Philip Hershkovitz, Assistant Curator of Mammals in Chicago Natural History Museum, extends the ranges of several Colombian species and includes three adult specimens of an undescribed race of *Tinamus osgoodi* Conover, 1949. This strikingly distinct species is an addition to the Colombian fauna, having been known previously only from the Marcapata Valley, Cuzco, Peru, some 1,200 miles to the southward. I take pleasure in naming the Colombian representative in honor of its collector.

### *Tinamus osgoodi hershkovitzi* subsp. nov.

*Type*.—Chicago Natural History Museum no. 217368, from San Adolfo, Río Aguas Claras, Acevedo, Huila, Colombia. Altitude 1,500 meters. Adult male collected July 1, 1951, by Philip Hershkovitz. Original number 101.

*Diagnosis*.—Similar to the nominate race of the Marcapata Valley, Cuzco, Peru, but wing coverts and back virtually concolor, and feathers of the under tail coverts rich cinnamon medially (not ochraceous buff), the edgings more or less tinged with dusky, but with little or no distinct speckling, vermiculation, or barring.

*Measurements*.—Male adult: wing 236; culmen 34; tarsus 75. Female adults: wing 258, 247; culmen 38, 35; tarsus 78, 75.

*Range*.—Known from only three specimens, collected at the type locality.

*Remarks*.—Traylor (1952, p. 18) has shown that *osgoodi* is most nearly related to *Tinamus tao*, which ranges in Brazil south of the Amazon to Mato Grosso and to the tropical zones of eastern Bolivia, Peru, Ecuador, British Guiana, northern Venezuela, and Colombia, and to *Tinamus major*, which ranges from Mexico through Central America to Peru, and Mato Grosso, Brazil. It agrees closely with

AL  
1953

*major* in wing length and with *tao* in tarsal length. These proportional relationships are corroborated by the measurements of *T. o. herskovitzi*.

## REFERENCES

CONOVER, BOARDMAN

1949. A new species of *Tinamus* from Peru. *Fieldiana, Zool.*, **31**: 263-266, figs. 66, 67.

TRAYLOR, MELVIN A., JR.

1952. Notes on birds from the Marcapata Valley, Peru. *Fieldiana, Zool.*, **34**: 17-23.





70.5

I

34 '18

nat. Hist.













UNIVERSITY OF ILLINOIS-URBANA

590 5F1 C001  
FIELDIANA ZOOLOGY\$CHGO  
34 1952-57



3 0112 009379659